

## Summary of adjustments to CPR algorithms in suspected or confirmed COVID-19 patients.

### Reduce provider exposure

- Don PPE before entering the room/scene
- Limit personnel
- Consider using mechanical CPR devices for adults and adolescents who meet height and weight criteria
- Communicate COVID-19 status to any new providers

### Prioritize oxygenation and ventilation strategies with lower aerosolization risk

- Use a HEPA filter, if available, for all ventilation
- Intubate early with a cuffed tube, if possible, and connect to mechanical ventilator, when able
- Engage the intubator with highest chance of first-pass success
- Pause chest compressions to intubate
- Consider use of video laryngoscopy, if available
- Before intubation, use a bag-mask device (or T-piece in neonates) with a HEPA filter and a tight seal
- For adults, consider passive oxygenation with nonrebreathing face mask as alternative to bag- mask device for short duration
- If intubation delayed, consider supraglottic airway
- Minimize closed circuit disconnections


### Consider resuscitation appropriateness

- Address goals of care
- Adopt policies to guide determination, taking into account patient risk factors for survival



# BLS Healthcare Provider Adult Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Updated April 2020




© 2020 American Heart Association



™ The heart and / icon on its own and the heart and / icon followed by another icon or words are trademarks of the Heart and Stroke Foundation of Canada.

# ACLS Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Updated April 2020



## CPR Quality

- Push hard (at least 5 cm [2 inches]) and fast (100-120/min) and allow complete chest recoil.
- Minimize interruptions in compressions.
- Avoid excessive ventilation.
- Change compressor every 2 minutes, or sooner if fatigued.
- If no advanced airway, 30:2 compression-ventilation ratio.
- Quantitative waveform capnography
  - If PETCO<sub>2</sub> <10 mm Hg, attempt to improve CPR quality.
- Intra-arterial pressure
  - If relaxation phase (diastolic) pressure <20 mm Hg, attempt to improve CPR quality.

## Shock Energy for Defibrillation

- Biphasic:** Manufacturer recommendation (eg, initial dose of 120-200 J); if unknown, use maximum available. Second and subsequent doses should be equivalent, and higher doses may be considered.
- Monophasic:** 360 J

## Advanced Airway

- Minimize closed-circuit disconnection
- Use intubator with highest likelihood of first pass success
- Consider video laryngoscopy
- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions

## Drug Therapy

- Epinephrine IV/IO dose:** 1 mg every 3-5 minutes
- Amiodarone IV/IO dose:** First dose: 300 mg bolus. Second dose: 150 mg. or **Lidocaine IV/IO dose:** First dose: 1-1.5 mg/kg. Second dose: 0.5-0.75 mg/kg.

## Return of Spontaneous Circulation (ROSC)


- Pulse and blood pressure
- Abrupt sustained increase in PETCO<sub>2</sub> (typically ≥40 mm Hg)
- Spontaneous arterial pressure waves with intra-arterial monitoring

## Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary

# BLS Healthcare Provider Pediatric Cardiac Arrest Algorithm for the Single Rescuer for Suspected or Confirmed COVID-19 Patients


Updated April 2020



© 2020 American Heart Association

# BLS Healthcare Provider Pediatric Cardiac Arrest Algorithm for 2 or More Rescuers for Suspected or Confirmed COVID-19 Patients


Updated April 2020



© 2020 American Heart Association

# Pediatric Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Updated April 2020



## CPR Quality

- Push hard ( $\geq 1/3$  of anteroposterior diameter of chest) and fast (100-120/min) and allow complete chest recoil.
- Minimize interruptions in compressions.
- Avoid excessive ventilation.
- Change compressor every 2 minutes, or sooner if fatigued.
- If no advanced airway, 15:2 compression-ventilation ratio.

## Shock Energy for Defibrillation

First shock 2 J/kg, second shock 4 J/kg, subsequent shocks  $\geq 4$  J/kg, maximum 10 J/kg or adult dose

## Advanced Airway

- **Minimize closed-circuit disconnection**
- **Use intubator with highest likelihood of first pass success**
- **Consider video laryngoscopy**
- **Prefer cuffed endotracheal tube if available**
- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions

## Drug Therapy

- **Epinephrine IO/IV dose:** 0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration). Repeat every 3-5 minutes.
- **Amiodarone IO/IV dose:** 5 mg/kg bolus during cardiac arrest. May repeat up to 2 times for refractory VF/pulseless VT.
- **Lidocaine IO/IV dose:** Initial: 1 mg/kg loading dose. Maintenance: 20-50 mcg/kg per minute infusion (repeat bolus dose if infusion initiated >15 minutes after initial bolus therapy).

## Return of Spontaneous Circulation (ROSC)

- Pulse and blood pressure
- Spontaneous arterial pressure waves with intra-arterial monitoring

## Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypoglycemia
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary

© 2020 American Heart Association



™ The heart and / icon on its own and the heart and / icon followed by another icon or words are trademarks of the Heart and Stroke Foundation of Canada.